arXiv:1611.02083v1 [quant-ph] 7 Nov 2016

Perturbative treatment of the non-linear q-Schrödinger and q-Klein-Gordon equations

D. J. Zamora[1], M. C. Rocca[1], A. Plastino[1], and G. L. Ferri[2]

[1] La Plata National University and Argentina's National Research Council (IFLP-CCT-CONICET)-C. C. 727, 1900 La Plata - Argentina

[2] Faculty of Exact and Natural Sciences, La Pampa National University Uruguay 151, Santa Rosa, La Pampa, Argentina

November 8, 2016

Abstract

Interesting nonlinear generalization of both Schrödinger's and Klein-Gordon's equations have been recently advanced by Tsallis, Rego-Monteiro, and Tsallis (NRT) in [Phys. Rev. Lett. **106**, 140601 (2011)]. There is much current activity going on in this area. The non-linearity is governed by a real parameter q. It is a fact that the ensuing non linear q-Schrödinger and q-Klein-Gordon equations are natural manifestations of very high energy phenomena, as verified by LHC-experiments. This happens for $\mathsf{q}-$values close to unity [Nucl. Phys. A **955**, 16 (2016), Nucl. Phys. A **948**, 19 (2016)]. It is also well known that q-exponential behavior is found in quite different settings. An explanation for such phenomenon was given in [Physica A **388**, 601 (2009)] with reference to empirical scenarios in which data are collected via set-ups that effect a normalization plus data's pre-processing. Precisely, the ensuing normalized output was there shown to be q-exponentially distributed if the input data display elliptical symmetry, generalization of spherical symmetry, a frequent situation. This makes it difficult, for q-values close to unity, to ascertain whether one is dealing with solutions to the ordinary Schrödinger

equation (whose free particle solutions are exponentials, and for which $q = 1$) or with its NRT nonlinear q-generalizations, whose free particle solutions are q-exponentials. In this work we provide a careful analysis of the $q \sim 1$ instance via a perturbative analysis of the NRT equations.
Keywords: Non-linear Schrödinger equation; Non-linear Klein-Gordon equation; First order solution.

1 Introduction

Empirical data indicates that power-law behavior in the observed probability distribution of interesting quantities is quite frequent in the natural world [1]. It was shown in [2] that one reason for this phenomenon is detector-normalization. In most measurement devices one finds a pre-processing step that impedes that the device be overwhelmed by data of too large amplitude that might damage he hardware. One then appeals to statistically normalization of input data. The data are first centered by subtraction of their estimated mean, and then scaled with their estimated standard deviation. It is shown in [2] that the concomitant proceedings transform Gaussian input data into q-Gaussian output one. We remind that a q-exponential is defined as

$$e_q(x) = [1 + (1-q)x]^{\frac{1}{1-q}}; \;\; ()e_{q=1}(x) = \exp x). \tag{1.1}$$

In view of the above empirical considerations, it becomes clear that in the immediate neighborhood of $q = 1$ it is very difficult to ascertain whether one is dealing with exponentials or with q-exponentials. The first instance corresponds to free-particle solutions of the celebrated Schrödinger equation, while the second instance corresponds to free particle solutions to its q-nonlinear generalization advanced in [5] (see also [6, 7]), the so called NRT equation. If we confront a particle flow, which of the two equations is governing it, the linear or the nonlinear one?

In order to help finding an adequate answer we carefully study here perturbative NRT solutions in a $q \sim 1$ scenario. We hope that our considerations will shed some light on these matters.

1.1 Motivation

The motivation of this paper resides in the fact that both the q-Schrödinger equation and the q-Klein-Gordon equation are natural manifestations of very high energy phenomena [8, 9], as verified by LHC-experiments [10], for which q is close to unity. In such a case, the two equations mentioned above approach the customary Schrödinger and Klein-Gordon equations, becoming identical to them in the limit $q \to 1$. The q-Schrödinger equation is

$$i\hbar \frac{\partial}{\partial t}\left[\frac{\psi(\vec{x},t)}{\psi(0,0)}\right]^q + \frac{\hbar^2}{2m}\triangle\left[\frac{\psi(\vec{x},t)}{\psi(0,0)}\right] = 0, \tag{1.2}$$

and for q close to unity we are allowed to write

$$\left[\frac{\psi(\vec{x},t)}{\psi(0,0)}\right]^q = \frac{\psi(\vec{x},t)}{\psi(0,0)} + (q-1)\frac{\psi(\vec{x},t)}{\psi(0,0)}\ln\left[\frac{\psi(\vec{x},t)}{\psi(0,0)}\right]. \tag{1.3}$$

The second term on the right side of (1.3) is negligible so that one writes

$$i\hbar\frac{\partial}{\partial t}\left[\frac{\psi(\vec{x},t)}{\psi(0,0)}\right] + \frac{\hbar^2}{2m}\triangle\left[\frac{\psi(\vec{x},t)}{\psi(0,0)}\right] = 0, \tag{1.4}$$

or

$$i\hbar\frac{\partial}{\partial t}\psi(\vec{x},t) + \frac{\hbar^2}{2m}\triangle\psi(\vec{x},t) = 0, \tag{1.5}$$

the linear, conventional de Schrödinger equation.

An analogous situation arises for the q-Klein-Gordon equation. We have

$$\Box\left[\frac{\phi(\vec{x},t)}{\phi(0,0)}\right] + \frac{qm^2c^2}{\hbar^2}\left[\frac{\phi(\vec{x},t)}{\phi(0,0)}\right]^{2q-1} = 0. \tag{1.6}$$

For q close to unity one has

$$\left[\frac{\phi(\vec{x},t)}{\phi(0,0)}\right]^{2q-1} = \frac{\phi(\vec{x},t)}{\phi(0,0)} + 2(q-1)\frac{\phi(\vec{x},t)}{\phi(0,0)}\ln\left[\frac{\phi(\vec{x},t)}{\phi(0,0)}\right]. \tag{1.7}$$

Once more, the second term on the right is negligible. As a consequence, we can write

$$\Box\left[\frac{\phi(\vec{x},t)}{\phi(0,0)}\right] + \frac{qm^2c^2}{\hbar^2}\left[\frac{\phi(\vec{x},t)}{\phi(0,0)}\right] = 0, \tag{1.8}$$

or

$$\Box\phi(\vec{x},t) + \frac{qm^2c^2}{\hbar^2}\phi(\vec{x},t) = 0, \tag{1.9}$$

and then, since q is close to unity,

$$\Box\phi(\vec{x},t) + \frac{m^2c^2}{\hbar^2}\phi(\vec{x},t) = 0. \tag{1.10}$$

We see then that it is very important to obtain approximate q-solutions for the two above scenarios, since these are two instances that correspond to intermediate energies [11].

2 First order expansion of the q exponential as a solution of the non-linear NRT q-Schrödinger equation

As a first task we will prove that the first order Taylor expansion, around $q = 1$, of q-exponential function e_q, is a solution of to the non-linear q-Schrödinger equation advanced in [5]

$$i\hbar\frac{\partial}{\partial t}\left[\frac{\psi(x,t)}{\psi(0,0)}\right]^q = H_0\left[\frac{\psi(x,t)}{\psi(0,0)}\right] \tag{2.1}$$

$\psi(0,0)$ a fixed wave function, where

$$H_0 = \frac{p^2}{2m} = \frac{-\hbar^2}{2m}\nabla^2 \tag{2.2}$$

In our case: $\psi \sim e_q$, defined as

$$e_q = [1 + \frac{i}{\hbar}(1-q)(px - Et)]^{\frac{1}{1-q}} \tag{2.3}$$

Calling $z = \frac{i}{\hbar}(px - Et)$ we deal with

$$e_q = [1 + (1-q)z]^{\frac{1}{1-q}} \tag{2.4}$$

The function e_q tends to the usual exponential when $q \to 1$

$$e_q(z)|_{q\to 1} = e^z \tag{2.5}$$

We observe that $\psi(0,0) = 1$. Thus, equation (2.1) reduces to

$$i\hbar\frac{\partial\psi^q}{\partial t} = H_0\psi(x,t). \tag{2.6}$$

2.1 First order expansion of $\psi = e_q$

We obtain, after a somewhat lengthy manipulation (see Appendix A)

$$\psi \simeq e^z + (q-1)\frac{z^2}{2}e^z,$$

or

$$\psi \simeq e^z \left[1 + (q-1)\frac{z^2}{2}\right]. \tag{2.7}$$

Note that the last relation differs form the pure exponential, for q close to unity, just by the term $(q-1)x^2/2$ above.

Moreover, we have

$$\psi \simeq e^{\frac{i}{\hbar}(px-Et)} \left[1 + (1-q)\frac{(px-Et)^2}{2\hbar^2}\right] \tag{2.8}$$

We need now the expansion for the second derivative of e_q with respect to x. This involves again some extensive manipulation (see Appendix A) and one finds

$$\frac{\partial^2\psi}{\partial x^2} = -\frac{p^2}{\hbar^2} e^{\frac{i}{\hbar}(px-Et)} \left[q + \frac{2i}{\hbar}(q-1)(px-Et) - \frac{(q-1)}{2\hbar^2}(px-Et)^2\right]. \tag{2.9}$$

Obviously, one also needs the first order expansion of ψ^q. This expansion is in the variable q, around $q = 1$, and has no obvious quantum mechanics counterpart. One writes

$$\psi^q = e^{q \ln \psi}.$$

Thus,

$$\frac{\partial\psi^q}{\partial q} = \frac{\partial(q \ln\psi)}{\partial q}\psi^q = \left(\ln\psi + \frac{q}{\psi}\frac{\partial\psi}{\partial q}\right)\psi^q = \psi^q \ln\psi + q\psi^{q-1}\frac{\partial\psi}{\partial q}. \tag{2.10}$$

For q=1 we obtain

$$\left.\frac{\partial\psi^q}{\partial q}\right|_{q=1} = \psi|_{q=1}\,(\ln\psi)|_{q=1} + \left.\frac{\partial\psi}{\partial q}\right|_{q=1}.$$

Since we know that

- $\psi|_{q=1} = e^z$
- $(\ln\psi)|_{q=1} = z$
- $\left.\frac{\partial\psi}{\partial q}\right|_{q=1} = \frac{z^2}{2}e^z$,

then,

$$\left.\frac{\partial \psi^q}{\partial q}\right|_{q=1} = ze^z + \frac{z^2}{2}e^z, \tag{2.11}$$

$$\left.\frac{\partial \psi^q}{\partial q}\right|_{q=1} = \frac{i}{\hbar}(px - Et)e^{\frac{i}{\hbar}(px-Et)} - \frac{(px-Et)^2}{2\hbar^2}e^{\frac{i}{\hbar}(px-Et)}. \tag{2.12}$$

The first order expansion of ψ^q is then, up to a normalization constant

$$\psi^q \simeq \psi^q|_{q=1} + (q-1)\left.\frac{\partial \psi^q}{\partial q}\right|_{q=1}.$$

Replacing here equations (2.5) and (2.12) we have

$$\psi^q \simeq e^{\frac{i}{\hbar}(px-Et)} + (q-1)\left[\frac{i}{\hbar}(px-Et)e^{\frac{i}{\hbar}(px-Et)} - \frac{(px-Et)^2}{2\hbar^2}e^{\frac{i}{\hbar}(px-Et)}\right]. \tag{2.13}$$

Finally, we require the time derivative. The first order time-derivative of ψ^q is

$$\frac{\partial \psi^q}{\partial t} = \frac{-iE}{\hbar}e^{\frac{i}{\hbar}(px-Et)} + (q-1)\left[\frac{-iE}{\hbar}e^{\frac{i}{\hbar}(px-Et)} + \frac{2E}{\hbar^2}(px-Et)e^{\frac{i}{\hbar}(px-Et)}\right.$$

$$\left. + \frac{iE}{2\hbar^3}(px-Et)^2 e^{\frac{i}{\hbar}(px-Et)}\right]$$

, or

$$\frac{\partial \psi^q}{\partial t} = \frac{-iE}{\hbar}e^{\frac{i}{\hbar}(px-Et)}\left[q + \frac{2i(q-1)}{\hbar}(px-Et) - \frac{(q-1)}{2\hbar^2}(px-Et)^2\right]. \tag{2.14}$$

2.2 Solution to the non-linear q-Schrödinger equation

Replacing equations (2.9) and (2.14) into the non-linear q-Schrödinger equation (2.6), we find that the first order Taylor's expansion of a q-exponential is indeed a solution of this equation with the usual eigenvalue corresponding to the free particle, $E = p^2/2m$. We have thus achieved self consistency, which is the first result of the present communication.

2.3 Comparison between the exact and approximate solutions

In this subsection we intend making comparisons between the approximate and exact solutions of q-Schrödinger equation. To this end, we first try to evaluate the modulus of the ratio of the functions (2.8) and (2.3). (We call to this ratio R). As an example, we display four figures corresponding to an electron and a proton with 1 MeV of energy at $t = 0$ (Figures 1,2,3, and 4). Note that, for a range of x very large in terms of atomic or nuclear relevant distances, the ratio is essentially unity. Our approximation can then be deemed very good.

3 Separation of variables

We will see here that, for $\psi(x,t) = f(t)g(x)$, the equations for, respectively, f and g, keep the form of the nonlinear equation for ψ (2.1). From (2.1), the non-linear q-Schrödinger equations for separate variables [$\psi(t,x) = f(t)g(x)$] are given by [3]

$$i\hbar\frac{\partial f^q}{\partial t} = \lambda f(t), \tag{3.1}$$

$$\frac{-\hbar^2}{2m}\frac{\partial^2 g}{\partial x^2} = \lambda g^q. \tag{3.2}$$

We should now prove that the first order expansion around $q = 1$ of the functions

$$f(t) = \left[1 + \frac{i}{\hbar}\frac{(1-q)}{q}Et\right]^{\frac{1}{q-1}}, \tag{3.3}$$

$$g(x) = \left[1 + \frac{i}{\hbar}\frac{(1-q)}{\sqrt{2(q+1)}}px\right]^{\frac{2}{1-q}}, \tag{3.4}$$

are solutions of their respective differential equations. This is done in Supplementary Materials (SM) B.

3.1 Solution to the differential equations for f and g

From the Appendix B we obtain the two relations

$$f(t) \simeq e^{\frac{-iEt}{\hbar}} \left[1 + (q-1)\left(\frac{iEt}{\hbar} + \frac{E^2t^2}{2\hbar^2}\right)\right], \tag{3.5}$$

and

$$\frac{\partial f^q}{\partial t} = -\frac{iE}{\hbar} e^{\frac{-iEt}{\hbar}} \left[1 + (q-1)\frac{E^2t^2}{2\hbar^2} + (q-1)\frac{iEt}{\hbar}\right]. \tag{3.6}$$

1) Now, replacing equations (3.5) and (3.6) (see Appendix B) into (3.1), we observe the first order Taylor expansion of f is a solution of (3.1), with $\lambda = E = p^2/2m$. Again, self-consistency has been reached. We pass next to SM C to deal with $g(x)$.

2) From Appendix C we get

$$g^q \simeq e^{\frac{ipx}{\hbar}} \left[1 + \frac{3(q-1)}{4}\frac{ipx}{\hbar} - \frac{(q-1)}{4}\frac{p^2x^2}{\hbar^2}\right], \tag{3.7}$$

and

$$\frac{\partial^2 g}{\partial x^2} = -\frac{p^2}{\hbar^2} e^{\frac{ipx}{\hbar}} \left[1 - \frac{3(1-q)}{4}\frac{ipx}{\hbar} + \frac{(1-q)}{4}\frac{p^2x^2}{\hbar^2}\right]. \tag{3.8}$$

Replacing (3.7) and (3.8) of Appendix C into (3.2), we find that the first order Taylor's expansion is a solution of this last equation with $\lambda = E = p^2/2m$.

4 First order treatment of a q-Gaussian

As mentioned in the Introduction, this issue is of paramount importance. From [12], selecting $mq\alpha = 1$ in order to simplify the calculations, we have for the q-Gaussian:

$$\psi(x,t) = \left\{1 + (q-1)\left[a(t)x^2 + b(t)x + c(t)\right]\right\}^{\frac{1}{1-q}} \tag{4.1}$$

where

$$a(t) = \frac{mq}{1 + i\hbar(q+1)t} \tag{4.2}$$

$$b(t) = \frac{1}{\beta[1 + i\hbar(q+1)t]} \tag{4.3}$$

$$c(t) = \left(\frac{1}{q-1} - \frac{1}{4mq\beta^2}\right)[1 + i\hbar(q+1)t]^{\frac{q-1}{q+1}} + \frac{1}{4mq\beta^2[1 + i\hbar(q+1)t]} + \frac{1}{1-q} \tag{4.4}$$

Writing up to first order

$$\begin{aligned} a(t) &= a_1(t) + (q-1)a_2(t) \\ b(t) &= b_1(t) + (q-1)b_2(t) \\ c(t) &= c_1(t) + (q-1)c_2(t) \end{aligned} \tag{4.5}$$

we obtain from (4.2), (4.3), and (4.4):

$$a_1(t) = \frac{m}{1 + 2i\hbar t} \tag{4.6}$$

$$a_2(t) = \frac{m(1 + i\hbar t)}{(1 + 2i\hbar t)^2} \tag{4.7}$$

$$b_1(t) = \frac{1}{\beta(1 + 2i\hbar t)} \tag{4.8}$$

$$b_2(t) = -\frac{i\hbar t}{\beta(1 + 2i\hbar t)^2} \tag{4.9}$$

$$c_1(t) = \frac{1}{2}\ln(1 + 2i\hbar t) - \frac{i\hbar t}{2m\beta^2(1 + 2i\hbar t)} \tag{4.10}$$

$$c_2(t) = \frac{1}{4m\beta^2} + \frac{i\hbar t}{2(1 + 2i\hbar t)} - \frac{(1 + 3i\hbar t)}{4m\beta^2(1 + 2i\hbar t)^2} + \frac{1}{8}\ln^2(1 + 2i\hbar t) - \frac{1 + 2m\beta^2}{8m\beta^2}\ln(1 + 2i\hbar t) \tag{4.11}$$

The first order approximation for the q-Gaussian is now

$$\psi(x,t) = \left\{1 - (q-1)\left\{a_2(t)x^2 + b_2(t)x + c_2(t) - \frac{1}{2}\left[a_1(t)x^2 + b_1(t)xc_1(t)\right]^2\right\}\right\} \otimes e^{-[a_1(t)x^2 + b_1(t)x + c_1(t)]} \tag{4.12}$$

By construction $\psi(x,t)$, as given by (4.11), is a first order solution to (2.1). Fig. 5 displays the ratio between (4.11) and (2.1) versus distance x (in absolute units) for $1-q = 10^{-3}$. Remark that the ratio is essentially unity for distances very much larger than atomic or nuclear ones. The approximation is quite good then.

4.1 Comparison between the exact and approximate solutions

To have an idea about the quality of the first order treatment of the qGaussian ans the exact solution we evaluate the modulus of the ratio of eq. (4.12) and (4.1) in a semi logarithmic scale. This is given in Fig, 5.

5 Non-linear q-Klein-Gordon equation

We should prove that the development of e_q is a solution of the following equation:

$$\frac{1}{c^2}\frac{\partial^2 F}{\partial t^2} - \frac{\partial^2 F}{\partial x^2} + \frac{q m^2 c^2}{\hbar^2} F^{2q-1} = 0 \tag{5.1}$$

This equation was advanced in [5] and re-obtained in [4].

In our case $F = e_q$, which, let us remind the reader, is defined as

$$e_q = [1 + i(1-q)(kx - \omega t)]^{\frac{1}{1-q}} \tag{5.2}$$

By analogy to equations (2.8) and (2.9) we know the expansions of e_q and of its derivative with respect to x, respectively cast as

$$F \simeq e^{i(kx-\omega t)}\left[1 + (1-q)\frac{(kx-\omega t)^2}{2}\right], \tag{5.3}$$

$$\frac{\partial^2 F}{\partial x^2} = -k^2 e^{i(kx-\omega t)}\left[q + 2i(q-1)(kx-\omega t) - \frac{(q-1)}{2}(kx-\omega t)^2\right]. \tag{5.4}$$

We should now calculate the second derivative with respect to t and the first order expansion of qF^{2q-1}, and so on. This is done in Appendix D.

5.1 Solution to the Klein-Gordon equation

Replacing (D.8), (D.2), and (5.4) into (5.1) we find that the first order Taylor's expansion of the q-exponential is a solution of this last equation. Again, self-consistency has been achieved.

6 Conclusions

We have exhaustively analyzed a first order perturbation-treatment (in q) of both the nonlinear q-Schrödinger and q-Klein Gordon partial differential equations. We have shown that, for small values of $\mathsf{q}-\mathsf{1}$, the approximation is quite good. This is of physical relevance because, as discussed in [8, 9], these q values are the relevant ones in the range of energies of interest for intermediate and high energy physics.

References

[1] N. Boccara, *Modeling complex systems*, (Springer, Berlin, 2004).

[2] C. Vignat, A. Plastino, Physica A **388** (2009) 601.

[3] A. Plastino, M. C. Rocca: Phys. Let. A **379**, 2690 (2015).

[4] A. Plastino, M. C. Rocca, G. L. Ferri and D. J. Zamora: Nuc. Phys. A **955**, 16 (2016).

[5] F. D. Nobre, M. A. Rego-Monteiro and C. Tsallis: Phys. Rev. Lett. **106**, 140601 (2011).

[6] A.R. Plastino, A.M.C. Souza, F.D. Nobre, C. Tsallis, Phys. Rev. A 90 (2014) 062134.

[7] A.R. Plastino, C. Tsallis, J. Math. Phys. 54 (2013) 041505; S. Curilef, A.R. Plastino, A. Plastino, Physica A 392 (2013) 2631; I.V. Toranzo, A.R. Plastino, J.S. Dehesa, A. Plastino, Physica A 392 (2013) 3945.

[8] A. Plastino, M. C. Rocca: Nucl. Phys. A **948** (2016) 19.

[9] A. Plastino, M. C. Rocca, G. L. Ferri, D. J. Zamora, Nucl. Phys. A 955 (2016) 1626.

[10] C. Tsallis, Introduction to Nonextensive Statistical Mechanics, Springer, Berlin, 2009; F. Barile, et al., ALICE Collaboration, EPJ Web Conf. 60 (2013) 13012; B. Abelev, et al., ALICE Collaboration, Phys. Rev. Lett. 111 (2013) 222301; Yu.V. Kharlov, ALICE Collaboration, Phys. At. Nucl. 76 (2013) 1497; ALICE Collaboration, Phys. Rev. C 91 (2015) 024609; ATLAS Collaboration, New J. Phys. 13 (2011) 053033; CMS Collaboration, J. High Energy Phys. 05 (2011) 064; CMS Collaboration, Eur. Phys. J. C 74 (2014) 2847; A. Adare, et al., PHENIX Collaboration, Phys. Rev. D 83 (2011) 052004; PHENIX Collaboration, Phys. Rev. C 83 (2011) 024909; PHENIX Collaboration, Phys. Rev. C 83 (2011) 064903; PHENIX Collaboration, Phys. Rev. C 84 (2011) 044902.

[11] A. Plastino, M. C. Rocca, G. L. Ferri and D. J. Zamora: Nuc. Phys. A **955**, 16 (2016).

[12] A. Plastino, M.C.Rocca: Physica A. **450**, 435 (2016).



Figure 1: Ratio R vs. x (in meters) for 1 MeV electrons and $q - 1 = 10^{-9}$.



Figure 2: Ratio R vs. x (in meters) for 1 MeV electrons and $q - 1 = 10^{-12}$.



Figure 3: Ratio R vs. x (in meters) for 1 MeV protons and $q - 1 = 10^{-9}$.



Figure 4: Ratio R vs. x (in meters) for 1 MeV protons and $q - 1 = 10^{-12}$.



Figure 5: q-Gaussian ratio of Eq. (4.12) over Eq. (4.1) vs. x (in absolute units) for $\mathsf{q}-1=10^{-3}$.

Appendix A

First order expansion of $\psi = e_q$

We write

$$\psi = e^{\frac{1}{1-q}\ln[1+(1-q)z]},$$

so that

$$\frac{\partial\psi}{\partial q} = \frac{\partial\left\{\frac{1}{1-q}\ln\left[1+(1-q)z\right]\right\}}{\partial q}\psi, \tag{A.1}$$

$$\frac{\partial\psi}{\partial q} = \left\{\frac{1}{(1-q)^2}\ln\left[1+(1-q)z\right] - \frac{z}{(1-q)\left[1+(1-q)z\right]}\right\}\psi.$$

As $\ln(1+x) = x - \frac{x^2}{2} + \cdots$, we obtain

$$\frac{\partial\psi}{\partial q} = \left\{\frac{1}{(1-q)^2}\left[(1-q)z - \frac{(1-q)^2z^2}{2}\right] - \frac{z}{(1-q)\left[1+(1-q)z\right]}\right\}\psi,$$

or

$$\frac{\partial\psi}{\partial q} = \left\{\frac{z}{(1-q)} - \frac{z}{(1-q)\left[1+(1-q)z\right]} - \frac{z^2}{2}\right\}\psi. \tag{A.2}$$

Let us evaluate $\frac{\partial\psi}{\partial q}$ at q=1:

$$\frac{\partial\psi}{\partial q} = \left\{z\left\{\frac{1}{(1-q)} - \frac{1}{(1-q)\left[1+(1-q)z\right]}\right\} - \frac{z^2}{2}\right\}\psi,$$

$$\frac{\partial\psi}{\partial q} = \left\{z\frac{\left[1+(1-q)z\right]-1}{(1-q)\left[1+(1-q)z\right]} - \frac{z^2}{2}\right\}\psi.$$

When $q \to 1$

$$\frac{\left[1+(1-q)z\right]-1}{(1-q)\left[1+(1-q)z\right]} \to \frac{0}{0}.$$

Using now L'Hopital's rule one has

$$\lim_{q\to 1}\frac{\partial\psi}{\partial q} = \lim_{q\to 1}\left\{\frac{z^2}{\left[1+(1-q)z\right]+(1-q)z} - \frac{z^2}{2}\right\}\psi,$$

$$\left.\frac{\partial\psi}{\partial q}\right|_{q=1} = \left(z^2 - \frac{z^2}{2}\right)e^z,$$

$$\left.\frac{\partial \psi}{\partial q}\right|_{q=1} = \frac{z^2}{2} e^z. \tag{A.3}$$

Thus, the first order Taylor's expansion of $\psi = e_q$ is

$$\psi \simeq \psi|_{q=1} + (q-1) \left.\frac{\partial \psi}{\partial q}\right|_{q=1}.$$

Second derivative with respect to x

The first order derivative respect to de variable x of e_q is

$$\frac{\partial \psi}{\partial x} = \frac{ip}{\hbar} e^{\frac{i}{\hbar}(px-Et)} - \frac{(q-1)}{2\hbar^2}\Big[2p(px-Et)e^{\frac{i}{\hbar}(px-Et)}$$

$$+\frac{ip}{\hbar}(px-Et)^2 e^{\frac{i}{\hbar}(px-Et)}\Big], \tag{A.4}$$

For the second derivative we have

$$\frac{\partial^2 \psi}{\partial x^2} = -\frac{p^2}{\hbar^2} e^{\frac{i}{\hbar}(px-Et)} - \frac{(q-1)}{2\hbar^2}\left[2p^2 e^{\frac{i}{\hbar}(px-Et)} + \frac{4ip^2}{\hbar}(px-Et)e^{\frac{i}{\hbar}(px-Et)}\right.$$

$$\left. -\frac{p^2}{\hbar^2}(px-Et)^2 e^{\frac{i}{\hbar}(px-Et)}\right],$$

or

$$\frac{\partial^2 \psi}{\partial x^2} = -\frac{p^2}{\hbar^2} e^{\frac{i}{\hbar}(px-Et)} - (q-1)\frac{p^2}{\hbar^2} e^{\frac{i}{\hbar}(px-Et)} -$$

$$(q-1)\frac{2ip^2}{\hbar^3}(px-Et)e^{\frac{i}{\hbar}(px-Et)} + (q-1)\frac{p^2}{2\hbar^4}(px-Et)^2 e^{\frac{i}{\hbar}(px-Et)}$$

.

Appendix B

First order expansion of f(t)

Since we can write

$$f(t) = \left[1 + \frac{i}{\hbar}\frac{(1-q)}{q}Et\right]^{\frac{1}{q-1}} = e^{\frac{1}{q-1}\ln\left[1+\frac{i}{\hbar}\frac{(1-q)}{q}Et\right]},$$

one has

$$\frac{\partial f}{\partial q} = \frac{\partial(\frac{1}{q-1}\ln\left[1+\frac{i}{\hbar}\frac{(1-q)}{q}Et\right])}{\partial q} f$$
$$= \left\{ -\frac{1}{(q-1)^2}\ln\left[1+\frac{i}{\hbar}\frac{(1-q)}{q}Et\right] + \right.$$
$$\left. \frac{1}{(q-1)\left[1+\frac{i}{\hbar}\frac{(1-q)}{q}Et\right]}\left[\frac{-iEt}{q\hbar} - \frac{(1-q)iEt}{q^2\hbar}\right] \right\} f.$$

Since $\ln(1+x) = x - \frac{x^2}{2} + \cdots$, we have

$$\frac{\partial f}{\partial q} = \left\{ -\frac{1}{(q-1)^2}\left[\frac{i}{\hbar}\frac{(1-q)}{q}Et + \frac{(1-q)^2}{2q^2}\frac{i}{\hbar}Et\right] \right.$$
$$\left. -\frac{1}{(q-1)\left[1+\frac{i}{\hbar}\frac{(1-q)}{q}Et\right]}\left[\frac{iEt}{q\hbar} + \frac{(1-q)}{q^2}\frac{iEt}{\hbar}\right] \right\} f. \qquad \text{(B.1)}$$

So as to evaluate f at $q = 1$ we rearrange terms in (B.1) and obtain

$$\frac{\partial f}{\partial q} = \left\{ \frac{iEt}{q\hbar}\left[-\frac{1}{(1-q)} + \frac{1}{(1-q)\left[1+\frac{i}{\hbar}\frac{(1-q)}{q}Et\right]} + \right.\right.$$
$$\left.\left. \frac{1}{q\left[1+\frac{i}{\hbar}\frac{(1-q)}{q}Et\right]}\right] - \frac{E^2t^2}{\hbar^2}\frac{1}{2q^2} \right\} f.$$

Moreover,

$$\frac{\partial f}{\partial q} = \left\{ \frac{iEt}{q\hbar}\left[\frac{1-\left[1+\frac{i}{\hbar}\frac{(1-q)}{q}Et\right]}{(1-q)\left[1+\frac{i}{\hbar}\frac{(1-q)}{q}Et\right]} + \frac{1}{q\left[1+\frac{i}{\hbar}\frac{(1-q)}{q}Et\right]}\right] - \frac{E^2t^2}{\hbar^2}\frac{1}{2q^2} \right\} f. \qquad \text{(B.2)}$$

As $q \to 1$

$$\frac{1-\left[1+\frac{i}{\hbar}\frac{(1-q)}{q}Et\right]}{(1-q)\left[1+\frac{i}{\hbar}\frac{(1-q)}{q}Et\right]} \to \frac{0}{0}.$$

Using again L'Hopital's rule we find

$$\lim_{q\to 1}\frac{1-\left[1+\frac{i}{\hbar}\frac{(1-q)}{q}Et\right]}{(1-q)\left[1+\frac{i}{\hbar}\frac{(1-q)}{q}Et\right]}=$$

$$\lim_{q\to 1}\frac{\frac{iEt}{q\hbar}+\frac{iEt(1-q)}{\hbar q^2}}{-\left[1+\frac{i}{\hbar}\frac{(1-q)}{q}Et\right]+(1-q)[\frac{iEt}{q\hbar}+\frac{iEt(1-q)}{\hbar q^2}}=-\frac{iEt}{\hbar}.$$

Thus,

$$\left.\frac{\partial f}{\partial q}\right|_{q=1}=\left\{\frac{iEt}{\hbar}\left[1-\frac{iEt}{\hbar}\right]-\frac{E^2t^2}{2\hbar^2}\right\}e^{\frac{-iEt}{\hbar}},$$

or

$$\left.\frac{\partial f}{\partial q}\right|_{q=1}=\left(\frac{iEt}{\hbar}+\frac{E^2t^2}{\hbar^2}-\frac{E^2t^2}{2\hbar^2}\right)e^{\frac{-iEt}{\hbar}}.$$

Moreover,

$$\left.\frac{\partial f}{\partial q}\right|_{q=1}=\left(\frac{iEt}{\hbar}+\frac{E^2t^2}{2\hbar^2}\right)e^{\frac{-iEt}{\hbar}}. \tag{B.3}$$

The first order Taylor's expansion of $f(t)$ is

$$f(t)\simeq f|_{q=1}+(q-1)\left.\frac{\partial f}{\partial q}\right|_{q=1}.$$

Replacing here (B.3) we have

$$f(t)\simeq e^{\frac{-iEt}{\hbar}}+(q-1)\left(\frac{iEt}{\hbar}+\frac{E^2t^2}{2\hbar^2}\right)e^{\frac{-iEt}{\hbar}},$$

or

$$f(t)\simeq e^{\frac{-iEt}{\hbar}}\left[1+(q-1)\left(\frac{iEt}{\hbar}+\frac{E^2t^2}{2\hbar^2}\right)\right]. \tag{B.4}$$

First order expansion of f^q

Following a similar procedure used above to obtain (2.12) we have

$$\left.\frac{\partial f^q}{\partial q}\right|_{q=1}=f|_{q=1}\ln f|_{q=1}+\left.\frac{\partial f}{\partial q}\right|_{q=1}. \tag{B.5}$$

We know that

- $f|_{q=1} = e^{\frac{-iEt}{\hbar}}$
- $(\ln f)|_{q=1} = -\frac{iEt}{\hbar}$
- $\left.\frac{\partial f}{\partial q}\right|_{q=1} = \left(\frac{iEt}{\hbar} + \frac{E^2t^2}{2\hbar^2}\right) e^{\frac{-iEt}{\hbar}}$.

Thus,

$$\left.\frac{\partial f^q}{\partial q}\right|_{q=1} = -\frac{iEt}{\hbar} e^{\frac{-iEt}{\hbar}} + e^{\frac{-iEt}{\hbar}} \left(\frac{iEt}{\hbar} + \frac{E^2t^2}{2\hbar^2}\right),$$

or

$$\left.\frac{\partial f^q}{\partial q}\right|_{q=1} = -\frac{iEt}{\hbar} e^{\frac{-iEt}{\hbar}} + \frac{iEt}{\hbar} e^{\frac{-iEt}{\hbar}} + \frac{E^2t^2}{2\hbar^2} e^{\frac{-iEt}{\hbar}}.$$

Moreover,

$$\left.\frac{\partial f^q}{\partial q}\right|_{q=1} = \frac{E^2t^2}{2\hbar^2} e^{\frac{-iEt}{\hbar}}. \tag{B.6}$$

The first order expansion of f^q is

$$f^q \simeq f^q|_{q=1} + (q-1) \left.\frac{\partial f^q}{\partial q}\right|_{q=1}.$$

Replacing here equation (B.6) we find

$$f^q \simeq e^{\frac{-iEt}{\hbar}} + (q-1)\frac{E^2t^2}{2\hbar^2} e^{\frac{-iEt}{\hbar}},$$

or, finally,

$$f^q \simeq e^{\frac{-iEt}{\hbar}} \left[1 + (q-1)\frac{E^2t^2}{2\hbar^2}\right]. \tag{B.7}$$

Derivative of f^q with respect to t

The first order derivative with respect to t of f^q is

$$\frac{\partial f^q}{\partial t} = -\frac{iE}{\hbar} e^{\frac{-iEt}{\hbar}} \left[1 + (q-1)\frac{E^2t^2}{2\hbar^2}\right] + e^{\frac{-iEt}{\hbar}} \frac{(q-1)E^2t}{\hbar^2},$$

or

$$\frac{\partial f^q}{\partial t} = -\frac{iE}{\hbar} e^{\frac{-iEt}{\hbar}} \left[1 + (q-1)\frac{E^2t^2}{2\hbar^2} + (q-1)\frac{iEt}{\hbar}\right]. \tag{B.8}$$

Appendix C

Expansion of g(x)

We can write

$$g(x) = \left[1 + \frac{i}{\hbar}\frac{(1-q)}{\sqrt{2(q+1)}}px\right]^{\frac{2}{1-q}} = e^{\frac{2}{1-q}\ln\left[1+\frac{i}{\hbar}\frac{(1-q)}{\sqrt{2(q+1)}}px\right]},$$

so that:

$$\begin{aligned}\frac{\partial g}{\partial q} &= \frac{\partial\left\{\frac{2}{1-q}\ln\left[1+\frac{i}{\hbar}\frac{(1-q)}{\sqrt{2(q+1)}}px\right]\right\}}{\partial q}g,\\ &= \left\{\frac{2}{(1-q)^2}\ln\left[1+\frac{i}{\hbar}\frac{(1-q)}{\sqrt{2(q+1)}}px\right] + \right.\\ &\frac{2}{(1-q)\left[1+\frac{i}{\hbar}\frac{(1-q)}{\sqrt{2(q+1)}}px\right]}\left[\frac{-ipx}{\hbar}\frac{1}{\sqrt{2(q+1)}}\right.\\ &\left.\left.-\frac{ipx}{\hbar}\frac{(1-q)}{[2(q+1)]^{3/2}}\right]\right\}g.\end{aligned}$$

Since $\ln(1+x) = x - \frac{x^2}{2} + \cdots$, we have

$$\begin{aligned}\frac{\partial g}{\partial q} = \left\{\frac{2}{(1-q)^2}\left[\frac{ipx}{\hbar}\frac{(1-q)}{\sqrt{2(q+1)}} + \frac{(1-q)^2}{2(q+1)}\frac{p^2x^2}{2\hbar^2}\right]\right.\\ \left.+\frac{2}{(1-q)\left[1+\frac{i}{\hbar}\frac{(1-q)}{\sqrt{2(q+1)}}px\right]}\left[-\frac{ipx}{\hbar}\frac{1}{\sqrt{2(q+1)}} - \frac{(1-q)}{[2(q+1)]^{3/2}}\frac{ipx}{\hbar}\right]\right\}g.\end{aligned} \tag{C.1}$$

So as to evaluate g at $q = 1$ we rearrange terms to obtain

$$\frac{\partial g}{\partial q} = \left\{\frac{ipx}{\hbar}\frac{2}{\sqrt{2(q+1)}(1-q)} + \frac{p^2x^2}{\hbar^2}\frac{1}{2(q+1)} -\right.$$

$$\frac{ipx}{\hbar}\frac{2}{\sqrt{2(q+1)}(1-q)\left[1+\frac{i}{\hbar}\frac{(1-q)}{\sqrt{2(q+1)}}px\right]}$$

$$-\frac{ipx}{\hbar}\frac{2}{[2(q+1)]^{3/2}\left[1+\frac{i}{\hbar}\frac{(1-q)}{\sqrt{2(q+1)}}px\right]}\Bigg\}g.$$

Moreover,

$$\frac{\partial g}{\partial q}=\left\{\frac{2ipx}{\hbar}\left[\frac{\left[1+\frac{i}{\hbar}\frac{(1-q)}{\sqrt{2(q+1)}}px\right]-1}{(1-q)\sqrt{2(q+1)}\left[1+\frac{i}{\hbar}\frac{(1-q)}{\sqrt{2(q+1)}}px\right]}-\right.\right.$$

$$\left.\left.\frac{1}{[2(q+1)]^{3/2}\left[1+\frac{i}{\hbar}\frac{(1-q)}{\sqrt{2(q+1)}}px\right]}\right]+\frac{p^2x^2}{\hbar^2}\frac{1}{2(q+1)}\right\}g.$$

As $q\to 1$,

$$\frac{\left[1+\frac{i}{\hbar}\frac{(1-q)}{\sqrt{2(q+1)}}px\right]-1}{(1-q)\sqrt{2(q+1)}\left[1+\frac{i}{\hbar}\frac{(1-q)}{\sqrt{2(q+1)}}px\right]}\to\frac{0}{0}. \tag{C.2}$$

Appealing once again to L'Hopital's rule we find

$$\lim_{q\to 1}\frac{\left[1+\frac{i}{\hbar}\frac{(1-q)}{\sqrt{2(q+1)}}px\right]-1}{(1-q)\sqrt{2(q+1)}\left[1+\frac{i}{\hbar}\frac{(1-q)}{\sqrt{2(q+1)}}px\right]}=$$

$$\lim_{q\to 1}\frac{\frac{-ipx}{\hbar}\frac{1}{\sqrt{2(q+1)}}-\frac{ipx}{\hbar}\frac{(1-q)}{[2(q+1)]^{3/2}}}{-\sqrt{2(q+1)}\left[1+\frac{i}{\hbar}\frac{(1-q)}{\sqrt{2(q+1)}}px\right]+(1-q)X},$$

$$=\frac{ipx}{4\hbar},$$

where

$$X = \frac{\left[1 + \frac{i}{\hbar}\frac{(1-q)}{\sqrt{2(q+1)}}px\right]}{\sqrt{2(q+1)}} + \sqrt{2(q+1)}\left[\frac{-ipx}{\hbar\sqrt{2(q+1)}} - \frac{(1-q)ipx}{\hbar[2(q+1)]^{3/2}}\right].$$

Thus,

$$\left.\frac{\partial g}{\partial q}\right|_{q=1} = \left\{\frac{2ipx}{\hbar}\left[\frac{ipx}{4\hbar} - \frac{1}{8}\right] + \frac{p^2x^2}{4\hbar^2}\right\} e^{\frac{ipx}{\hbar}},$$

or

$$\left.\frac{\partial g}{\partial q}\right|_{q=1} = \left(-\frac{ipx}{4\hbar} - \frac{p^2x^2}{2\hbar^2} + \frac{p^2x^2}{4\hbar^2}\right) e^{\frac{ipx}{\hbar}},$$

and, finally,

$$\left.\frac{\partial g}{\partial q}\right|_{q=1} = -\frac{1}{4}\left(\frac{ipx}{\hbar} + \frac{p^2x^2}{\hbar^2}\right) e^{\frac{ipx}{\hbar}}. \tag{C.3}$$

The first order expansion of $g(x)$ is

$$g(x) \simeq g|_{q=1} + (q-1)\left.\frac{\partial g}{\partial q}\right|_{q=1}.$$

Replacing here (C.3) we get

$$g(x) \simeq e^{\frac{ipx}{\hbar}} + \frac{(1-q)}{4}\left(\frac{ipx}{\hbar} + \frac{p^2x^2}{\hbar^2}\right) e^{\frac{ipx}{\hbar}},$$

or,

$$g(x) \simeq e^{\frac{ipx}{\hbar}}\left[1 + \frac{(1-q)}{4}\left(\frac{ipx}{\hbar} + \frac{p^2x^2}{\hbar^2}\right)\right]. \tag{C.4}$$

Second derivative of g with respect to x

The first order derivative of g is

$$\frac{\partial g}{\partial x} = \frac{ip}{\hbar}e^{\frac{ipx}{\hbar}}\left[1 + \frac{(1-q)}{4}\left(\frac{ipx}{\hbar} + \frac{p^2x^2}{\hbar^2}\right)\right] + e^{\frac{ipx}{\hbar}}\frac{(1-q)}{4}\left(\frac{ip}{\hbar} + \frac{2xp^2}{\hbar^2}\right). \tag{C.5}$$

For the second order derivative we have

$$\frac{\partial^2 g}{\partial x^2} = \frac{-p^2}{\hbar^2}e^{\frac{ipx}{\hbar}}\left[1 + \frac{(1-q)}{4}\left(\frac{ipx}{\hbar} + \frac{p^2x^2}{\hbar^2}\right)\right] + \frac{ip}{\hbar}e^{\frac{ipx}{\hbar}}\frac{(1-q)}{4}\left(\frac{ip}{\hbar} + \frac{2xp^2}{\hbar^2}\right)$$

$$+\frac{ip}{\hbar}e^{\frac{ipx}{\hbar}}\frac{(1-q)}{4}\left(\frac{ip}{\hbar}+\frac{2xp^2}{\hbar^2}\right)+e^{\frac{ipx}{\hbar}}\frac{(1-q)}{4}\frac{2p^2}{\hbar^2},$$

or

$$\frac{\partial^2 g}{\partial x^2}=-\frac{p^2}{\hbar^2}e^{\frac{ipx}{\hbar}}\left[1+\frac{(1-q)}{4}\frac{ipx}{\hbar}+\frac{(1-q)}{4}\frac{p^2x^2}{\hbar^2}-(1-q)\frac{ipx}{\hbar}\right],$$

so that, finally,

$$\frac{\partial^2 g}{\partial x^2}=-\frac{p^2}{\hbar^2}e^{\frac{ipx}{\hbar}}\left[1-\frac{3(1-q)}{4}\frac{ipx}{\hbar}+\frac{(1-q)}{4}\frac{p^2x^2}{\hbar^2}\right]. \tag{C.6}$$

First order expansion of g^q

Following a similar procedure to that used to obtain [1.5] in the paper we have now

$$\left.\frac{\partial g^q}{\partial q}\right|_{q=1}=\left.g\right|_{q=1}\ln\left.g\right|_{q=1}+\left.\frac{\partial g}{\partial q}\right|_{q=1}. \tag{C.7}$$

We know that

- $\left.g\right|_{q=1}=e^{\frac{ipx}{\hbar}}$
- $\left.(\ln g)\right|_{q=1}=\frac{ipx}{\hbar}$
- $\left.\frac{\partial g}{\partial q}\right|_{q=1}=-\frac{1}{4}\left(\frac{ipx}{\hbar}+\frac{p^2x^2}{\hbar^2}\right)e^{\frac{ipx}{\hbar}}$.

Thus,

$$\left.\frac{\partial g^q}{\partial q}\right|_{q=1}=\frac{ipx}{\hbar}e^{\frac{ipx}{\hbar}}-\frac{1}{4}e^{\frac{ipx}{\hbar}}\left(\frac{ipx}{\hbar}+\frac{p^2x^2}{\hbar^2}\right),$$

or,

$$\left.\frac{\partial g^q}{\partial q}\right|_{q=1}=\frac{ipx}{\hbar}e^{\frac{-iEt}{\hbar}}\left(1-\frac{1}{4}+\frac{ipx}{4\hbar}\right).$$

Moreover:

$$\left.\frac{\partial g^q}{\partial q}\right|_{q=1}=\frac{ipx}{\hbar}e^{\frac{ipx}{\hbar}}\left(\frac{3}{4}+\frac{ipx}{4\hbar}\right). \tag{C.8}$$

The first order expansion of g^q is

$$g^q\simeq\left.g^q\right|_{q=1}+(q-1)\left.\frac{\partial g^q}{\partial q}\right|_{q=1}. \tag{C.9}$$

Replacing Eq. (C.8) in this expansion we obtain

$$g^q \simeq e^{\frac{ipx}{\hbar}} + (q-1)\frac{ipx}{\hbar}e^{\frac{ipx}{\hbar}}\left(\frac{3}{4}+\frac{ipx}{4\hbar}\right),$$

or

$$g^q \simeq e^{\frac{ipx}{\hbar}}\left[1+\frac{3(q-1)}{4}\frac{ipx}{\hbar}-\frac{(q-1)}{4}\frac{p^2x^2}{\hbar^2}\right]. \tag{C.10}$$

Appendix D

Second derivative with respect to t

The first order derivative of F is

$$\frac{\partial F}{\partial t} = -i\omega e^{i(kx-\omega t)}\left[1+\frac{(1-q)}{2}(kx-\omega t)^2\right] - \omega e^{i(kx-\omega t)}(1-q)(kx-\omega t),$$

or

$$\frac{\partial F}{\partial t} = -i\omega e^{i(kx-\omega t)}\left[1-(1-q)i(kx-\omega t)+\frac{(1-q)}{2}(kx-\omega t)^2\right]. \tag{D.1}$$

For the second order derivative we have

$$\frac{\partial^2 F}{\partial t^2} = -\omega^2 e^{i(kx-\omega t)}\left[1-(1-q)i(kx-\omega t)+\frac{(1-q)}{2}(kx-\omega t)^2\right]$$

$$-i\omega e^{i(kx-\omega t)}\left[i\omega(1-q)-\omega(1-q)(kx-\omega t)\right],$$

or

$$\frac{\partial^2 F}{\partial t^2} = -\omega^2 e^{i(kx-\omega t)}\left[1-(1-q)i(kx-\omega t)+\right.$$

$$\left.\frac{(1-q)}{2}(kx-\omega t)^2-(1-q)-(1-q)i(kx-\omega t)\right],$$

so that, finally,

$$\frac{\partial^2 F}{\partial t^2} = -\omega^2 e^{i(kx-\omega t)}\left[q+2i(q-1)(kx-\omega t)-\frac{(q-1)}{2}(kx-\omega t)^2\right]. \tag{D.2}$$

First order expansion of qF^{2q-1}

The derivative of qF^{2q-1} with respect to q is

$$\frac{\partial(qF^{2q-1})}{\partial q} = F^{2q-1} + q\frac{\partial F^{2q-1}}{\partial q}. \tag{D.3}$$

We can write

$$F^{2q-1} = e^{(2q-1)\ln F},$$

so that

$$\frac{\partial F^{2q-1}}{\partial q} = \left[2\ln F + \frac{(2q-1)}{F}\frac{\partial F}{\partial q}\right] F^{2q-1},$$

$$\frac{\partial F^{2q-1}}{\partial q} = 2F^{2q-1}\ln F + (2q-1)F^{2q-2}\frac{\partial F}{\partial q}. \tag{D.4}$$

For the derivative of qF^{2q-1} we have

$$\frac{\partial(qF^{2q-1})}{\partial q} = F^{2q-1} + 2qF^{2q-1}\ln F + q(2q-1)F^{2q-2}\frac{\partial F}{\partial q}. \tag{D.5}$$

At $q = 1$ we obtain

$$\left.\frac{\partial(qF^{2q-1})}{\partial q}\right|_{q=1} = F|_{q=1} + 2\,F|_{q=1}\,\ln F|_{q=1} + \left.\frac{\partial F}{\partial q}\right|_{q=1}.$$

We know that

- $F|_{q=1} = e^{i(kx-\omega t)}$
- $(\ln F)|_{q=1} = i(kx-\omega t)$
- $\left.\frac{\partial F}{\partial q}\right|_{q=1} = -\frac{(kx-\omega t)^2}{2}e^{i(kx-\omega t)}$.

Thus,

$$\left.\frac{\partial(qF^{2q-1})}{\partial q}\right|_{q=1} = e^{i(kx-\omega t)} + 2i(kx-\omega t)e^{i(kx-\omega t)} - \frac{(kx-\omega t)^2}{2}e^{i(kx-\omega t)},$$

or

$$\left.\frac{\partial(qF^{2q-1})}{\partial q}\right|_{q=1} = e^{i(kx-\omega t)}\left[1 + 2i(kx-\omega t) - \frac{(kx-\omega t)^2}{2}\right]. \tag{D.6}$$

The first order expansion of qF^{2q-1} is

$$qF^{2q-1} \simeq (qF^{2q-1})\big|_{q=1} + (q-1)\left.\frac{\partial(qF^{2q-1})}{\partial q}\right|_{q=1}. \tag{D.7}$$

Replacing Eq. (D.6) in this expansion we obtain

$$qF^{2q-1} \simeq e^{i(kx-\omega t)} + (q-1)e^{i(kx-\omega t)}\left[1 + 2i(kx-\omega t) - \frac{(kx-\omega t)^2}{2}\right],$$

or

$$qF^{2q-1} \simeq e^{i(kx-\omega t)}\left[q + 2i(q-1)(kx-\omega t) - (q-1)\frac{(kx-\omega t)^2}{2}\right]. \tag{D.8}$$